

ON COMPUTABILITY OF JULIA SETS: ANSWERS TO QUESTIONS OF MILNOR AND SHUB

M. BRAVERMAN AND M. YAMPOLSKY

ABSTRACT. In this note we give answers to questions posed to us by J. Milnor and M. Shub, which shed further light on the structure of non-computable Julia sets.

1. INTRODUCTION

Computability of real sets. The reader is directed to [BY] for a more detailed discussion of the notion of computability of subsets of $\mathbb{R}^n$ as applied, in particular, to Julia sets. We recall the principal definitions here. The exposition below uses the concept of a *Turing Machine*. This is a standard model for a computer program employed by computer scientists. Readers unfamiliar with this concept should think instead of an algorithm written in their favorite programming language. These concepts are known to be equivalent.

Denote by $\mathbb{D}$ the set of the *dyadic rationals*, that is, rationals of the form $\frac{p}{2^m}$. We say that $\phi : \mathbb{N} \rightarrow \mathbb{D}$ is an *oracle* for a real number x , if $|x - \phi(n)| < 2^{-n}$ for all $n \in \mathbb{N}$. In other words, ϕ provides a good dyadic approximation for x . We say that a Turing Machine (further abbreviated as TM) M^ϕ is an *oracle machine*, if at every step of the computation M is allowed to query the value $\phi(n)$ for any n . This definition allows us to define the computability of real functions on compact sets.

Definition 1.1. We say that a function $f : [a, b] \rightarrow [c, d]$ is computable, if there exists an oracle TM $M^\phi(m)$ such that if ϕ is an oracle for $x \in [a, b]$, then on input m , M^ϕ outputs a $y \in \mathbb{D}$ such that $|y - f(x)| < 2^{-m}$.

It is worthwhile to note why the oracle mechanism is introduced. There are only countably many possible algorithms, and consequently only countably many *computable* real numbers which such algorithms can encode. Therefore, one wants to separate the hardness of encoding the real number x from the hardness of computing the value of the function $f(x)$, having the access to the value of x .

Let $K \subset \mathbb{R}^k$ be a compact set. We say that a TM M computes the set K if it approximates K in the *Hausdorff metric*. Recall that the Hausdorff metric is a metric on compact subsets of $\mathbb{R}^n$ defined by

$$(1) \quad d_H(X, Y) = \inf\{\epsilon > 0 \mid X \subset U_\epsilon(Y) \text{ and } Y \subset U_\epsilon(X)\},$$

where $U_\epsilon(S)$ is defined as the union of the set of ϵ -balls with centers in S .

We introduce a class $\mathcal{C}$ of sets which is dense in metric d_H among the compact sets and which has a natural correspondence to binary strings. Namely $\mathcal{C}$ is the set of finite unions

of dyadic balls:

$$\mathcal{C} = \left\{ \bigcup_{i=1}^n \overline{B(d_i, r_i)} \mid \text{where } d_i, r_i \in \mathbb{D} \right\}.$$

Members of $\mathcal{C}$ can be encoded as binary strings in a natural way.

We now define the notion of computability of subsets of $\mathbb{R}^n$ (see [Wei], and also [RW]).

Definition 1.2. We say that a compact set $K \subset \mathbb{R}^k$ is computable, if there exists a TM $M(d, n)$, where $d \in \mathbb{D}$, $n \in \mathbb{N}$ which outputs a value 1 if $\text{dist}(d, K) < 2^{-n}$, the value 0 if $\text{dist}(d, K) > 2 \cdot 2^{-n}$, and in the “in-between” case it halts and outputs either 0 or 1.

In other words, it computes, in the classical sense, a function from the family $\mathcal{F}_K$ of functions of the form

$$(2) \quad f(d, n) = \begin{cases} 0, & \text{if } \text{dist}(d, K) > 2 \cdot 2^{-n} \\ 1, & \text{if } \text{dist}(d, K) < 2^{-n} \\ 0 \text{ or } 1, & \text{otherwise} \end{cases}$$

Theorem 1.1. *For a compact $K \subset \mathbb{R}^k$ the following are equivalent:*

- (1) K is computable as per definition 1.2,
- (2) there exists a TM $M(m)$, such that on input m , $M(m)$ outputs an encoding of $C_m \in \mathcal{C}$ such that $d_H(K, C_m) < 2^{-m}$ (global computability),
- (3) the distance function $d_K(x) = \inf\{|x - y| \mid y \in K\}$ is computable as per definition 1.1.

Note that in the case $k = 2$ computability means that K can be drawn on a computer screen with arbitrarily good precision (if we imagine the screen as a lattice of pixels).

In the present paper we are interested in questions concerning the computability of the Julia set $J_c = J(f_c) = J(z^2 + c)$. Since there are uncountably many possible parameter values for c , we cannot expect for each c to have a machine M such that M computes J_c (recall that there are countably many TMs). On the other hand, it is reasonable to want M to compute J_c with an oracle access to c . Define the function $J : \mathbb{C} \rightarrow K^*$ (K^* is the set of all compact subsets of $\mathbb{C}$) by $J(c) = J(f_c)$. In a complete analogy to Definition 1.1 we can define

Definition 1.3. We say that a function $\kappa : S \rightarrow K^*$ for some bounded set S is computable, if there exists an oracle TM $M^\phi(d, n)$, where ϕ is an oracle for $x \in S$, which computes a function (2) of the family $\mathcal{F}_{\kappa(x)}$.

Equivalently, there exists an oracle TM $M^\phi(m)$ with ϕ again representing $x \in S$ such that on input m , M^ϕ outputs a $C \in \mathcal{C}$ such that $d_H(C, \kappa(x)) < 2^{-m}$.

In the case of Julia sets:

Definition 1.4. We say that J_c is computable if the function $J : d \mapsto J_d$ is computable on the set $\{c\}$.

We have the following (see [BBY2]):

Theorem 1.2. *Suppose that a TM M^ϕ computes the function J on a set $S \subset \mathbb{C}$. Then J is continuous on S in Hausdorff sense.*

Previous results. We have demonstrated in [BY]:

Theorem 1.3. *There exists a parameter value $c \in \mathbb{C}$ such that the Julia set of the quadratic polynomial $f_c(z) = z^2 + c$ is not computable.*

The quadratic polynomials in Theorem 1.3 possess Siegel disks. It was further shown by I. Binder and the authors of the present paper in [BBY1] that the absence of rotation domains, that is either Siegel disks or Herman rings, guarantees computability of the rational Julia set. This implies, in particular, that all Cremer quadratic Julia sets are computable – this despite the fact that no informative high resolution images of such sets have ever been produced. One expects, however, that such “bad” but still computable examples have high algorithmic complexity, which makes the computational cost of producing such a picture prohibitively high.

Two questions on computability of Julia sets. J. Milnor has asked us the following natural question:

Is the filled Julia set of a quadratic polynomial always computable?

In this paper we answer in the affirmative:

Theorem 1.4. *For any polynomial $p(z)$ there is an oracle Turing Machine $M^\phi(n)$ that given an oracle access to the coefficients of $p(z)$ and n , outputs a 2^{-n} -approximation of the filled Julia set $K_{p(z)}$.*

Moreover, in the case when $p(z) = z^2 + c$ is quadratic, only two machines suffice to compute all non-parabolic Julia sets: one for $c \in M$, and one for $c \notin M$.

This may come as a surprise, given the negative result for Julia sets. To gain some insight into how non-computability can be destroyed by filling in, consider the following toy example. Let $A : \mathbb{N} \rightarrow \{0, 1\}$ be any uncomputable predicate. Consider the set

$$\Omega_t = \begin{cases} S^1 \bigcup_{k \in \mathbb{N}, d_k=1} \{re^{2\pi i/k} \mid r \in [1 - \frac{1}{k}, 1]\} & \text{for } t = (0.d_1d_2d_3\dots)_2 \in [0, 1) \\ S^1 \bigcup_{k \in \mathbb{N}, A(k)=1} \{re^{2\pi i/k} \mid r \in [1 - \frac{1}{k}, 1]\} & \text{for } t = 1 \end{cases}$$

To avoid ambiguity, we always take the finite expansion for dyadic t ’s. An example of a set Ω_t is depicted on Figure 1. Firstly, note that if $t \in (0, 1)$ is not a computable real, then the set Ω_t is non-computable by a TM *without* an oracle for t . Moreover, even for a TM M^ϕ equipped with an oracle input for t , the set Ω_1 is clearly non-computable. However, when filled, every Ω_t becomes a computable set – the unit disk.

The question of M. Shub again has to do with fragility of non-computability. This time, instead of filling in a non-computable Julia set, we will make a “fuzzy” picture of it by letting the parameter c vary in a neighborhood.

To formalize this, consider the following definition. Let $\mathbb{J}$ be the subset of $\mathbb{C} \times \mathbb{C}$ given by

$$\mathbb{J} = \overline{\{(z, c) : z \in J_c\}}.$$

Shub has asked us:

Is the set $\mathbb{J}$ computable?

The answer again is “yes”:




FIGURE 1. Part of the picture of Ω_t for $t = (0.10101\dots)_2$

Theorem 1.5. *Let $d > 0$ be any computable real. Then the set*

$$\mathbb{J} \cap \mathbb{C} \times \overline{B(0, d)}$$

is a computable subset of $\mathbb{C} \times \mathbb{C}$.

Informally, we may think of projection of $\mathbb{J} \cap \mathbb{C} \times (c - \epsilon, c + \epsilon)$ to the first coordinate as the picture that a computer would produce when J_c itself is uncomputable.

To understand how the mechanism of non-computability is destroyed in this case, consider again the set Ω_t for $t \in (0, 1]$ as the toy model. The set $W = \overline{\{(z, t) : z \in \Omega_t, t \in (0, 1]\}} \subset \mathbb{C} \times \mathbb{R}$ is computable even though Ω_t itself is non-computable for $t = 1$. This happens because in the closure of W the “slice” corresponding to $t = 1$ is

$$S^1 \bigcup_{k \in \mathbb{N}} \{re^{2\pi i/k} \mid r \in [1 - \frac{1}{k}, 1]\} \supset \Omega_1.$$

This set “masks” the computational hardness of Ω_1 , and makes W computable.

2. COMPUTABILITY OF FILLED JULIA SETS

Our goal in this section is to prove Theorem 1.4. For a given polynomial $p(z)$ we construct a machine computing the corresponding filled Julia set K_p . We will use the following combinatorial information about p in the construction. Note that this information can be encoded using a finite number of bits.

- Information that would allow us to compute the non-repelling orbits of the polynomial with an arbitrary precision. Note that there are at most $\deg p - 1$ of them. Such information could, for example, consist of the list of periods k_i of such orbits; and for each i a finite collection of dyadic balls $\{D_i^j\}_{j=1}^{k_i}$ separating the points of the corresponding orbit from the other solutions of the equation $p^{k_i}(z) = z$. This allows for an arbitrarily precise approximation of the orbits by using an iterative root-finding algorithm for $p^{k_i}(z) = z$ in D_i^j .
- In the case of a hyperbolic or a parabolic orbit ζ , a domain of attraction D_ζ such that every orbit converging to ζ eventually reaches and stays in D_ζ . In the hyperbolic case D_ζ is just a collection of discs. In the parabolic case, it is a collection of sectors around the points of ζ , and it can be computed with an arbitrarily high precision.

- In the case of a Siegel disc D , information that would allow us to identify a repelling orbit ζ_D in the same connected component of K_p as D . Such an orbit always exists, and can be identified using a finite amount of combinatorial information.

2.1. Computing K_p . We are given a dyadic point $d \in \mathbb{D}$ and an $n \in \mathbb{N}$. Our goal is to always terminate and output 1 if $B(d, 2^{-n}) \cap K_p \neq \emptyset$ and to output 0 if $B(d, 2 \cdot 2^{-n}) \cap K_p = \emptyset$. We do it by constructing five machines. They are guaranteed to terminate each on a different condition, always with a valid answer. Together they cover all the possible cases.

Lemma 2.1. *There are five oracle machines M_{ext} , M_{jul} , M_{hyp} , M_{par} , M_{sieg} such that*

- (1) *if d is at distance $\geq \frac{4}{3} \cdot 2^{-n}$ from K_p , $M_{ext}(d, n)$ will halt and output 0. If d is at distance $\leq 2^{-n}$ from K_p , $M_{ext}(d, n)$ will never halt;*
- (2) *if d is at distance $\leq \frac{5}{3} \cdot 2^{-n}$ from J_p , $M_{jul}(d, n)$ will halt and output 1. If d is at distance $\geq 2 \cdot 2^{-n}$ from J_p , $M_{jul}(d, n)$ will never halt;*
- (3) *$M_{hyp}(d, n)$ halts and outputs 1 if and only if d is inside an attracting basin of a hyperbolic orbit of p ;*
- (4) *$M_{par}(d, n)$ halts and outputs 1 if and only if d is inside an attracting basin of a parabolic orbit of p ;*
- (5) *$M_{sieg}(d, n)$ halts and outputs 1 if the orbit of d reaches a Siegel disc, and d is at distance $\geq \frac{4}{3} \cdot 2^{-n}$ from J_p . It never halts if d is at distance $\geq 2 \cdot 2^{-n}$ from K_p .*

Recall the Fatou-Sullivan classification of Fatou components of a polynomial mapping of $\hat{\mathbb{C}}$ (see e.g. [Mil]):

Theorem 2.2 (Fatou-Sullivan classification). *Every Fatou component of a polynomial mapping of $\hat{\mathbb{C}}$ of degree at least two is a preimage of a periodic component. Every periodic component is of one of the following types: the immediate basin of an attracting (or a super-attracting) periodic point; a component of the immediate basin of a parabolic periodic point; a Siegel disk.*

Proof of Theorem 1.4, given Lemma 2.1. By Fatou-Sullivan classification it is not hard to see that for each (d, n) at least one of the machines halts. Moreover, by the definition of the machines, they always output a valid answer whenever they halt. Hence running the machines in parallel and returning the output of the first machine to halt gives the algorithm for computing K_p . $\square$

It remains to prove Lemma 2.1.

Proof. (of Lemma 2.1) We give a simple construction for each of the five machines.

- (1) M_{ext} : Take a large ball B such that $K_p \subset B$. Intuitively, we pull the ball back under p to get a good approximation of K_p . Let B_k be a $2^{-(n+3)}$ -approximation of the set $p^{-k}(B)$. Output 0 iff $B_k \cap B(d, \frac{7}{6} \cdot 2^{-n}) = \emptyset$. It is not hard to see that this algorithm satisfies the conditions on M_{ext} .
- (2) M_{jul} : Enumerate all the repelling periodic orbits of p . Let C_k be a $2^{-(n+3)}$ -approximation of the union of the first k orbits enumerated. Output 1 iff $d(d, C_k) < \frac{11}{6} \cdot 2^{-n}$. The repelling periodic orbits are all in J_p and are dense in this set. Hence the algorithm satisfies the conditions on M_{jul} .

- (3) M_{hyp} : Let z_k be a 2^{-k} -approximation of $p^k(d)$. If d is inside the basin of attraction for a some orbit ζ , then z_k for some k will be inside D_ζ for some k . Output 1 if z_k is at least 2^{-k} -far from the boundary of D_ζ .
- (4) M_{par} : Very similar to M_{hyp} . The only difference is that now we are checking for convergence to an attracting petal of a parabolic orbit.
- (5) M_{sieg} : This is the most interesting case. It is not hard to see that for each k , we can compute a union E_k of dyadic balls such that

$$\bigcup_{i=0}^k p^i \left(B(d, \frac{4}{3} \cdot 2^{-n}) \right) \subset E_k \subset \bigcup_{i=0}^k p^i \left(B(d, \frac{5}{3} \cdot 2^{-n}) \right).$$

Let c be the center of the Siegel disc (one of the centers, in case of an orbit), and let y be the given periodic point in the connected component of c . We terminate and output 1 if E_n separates c from y in $\mathbb{C}$ (or covers either one of them).

If d is inside the Siegel disc, then the forward images of $B(d, \frac{4}{3} \cdot 2^{-n})$ will cover an annulus in the disc that will separate c from the boundary of the disc, and in particular from y . Hence M_{sieg} will terminate and output 1.

If the distance from d to K_c is $\geq 2 \cdot 2^{-n}$, then $E_k \cap K_p = \emptyset$ for all k . In particular, E_k cannot separate c from y , since they are connected in K_p .

□

2.2. The quadratic case. In the quadratic case there is at most one non-repelling orbit. In the case there is a Siegel disc, the Julia set is connected, and any repelling periodic orbit can be taken as the orbit connected to the Siegel disc. In fact, it is not hard to see that if we exclude the parabolic case, one machine suffices to take care of all the *connected* filled Julia sets. As a corollary we get:

Corollary 2.3. *Denote by M the Mandelbrot set, and by P the set of c 's for which J_c is parabolic. The function $K : c \mapsto K_{z^2+c}$ is continuous in the Hausdorff metric on the set $M - P$.*

3. COMPUTABILITY OF THE SET $\mathbb{J}$

Recall that

$$\mathbb{J} = \overline{\{(z, c) : z \in J_c\}} \subset \mathbb{C}^2.$$

Theorem 1.5 asserts that $\mathbb{J}$ is computable. We prove it by showing that $\mathbb{J}$ is *weakly* computable.

Definition 3.1. We say that a set C is *weakly* computable if there is an oracle Turing Machine $M^\phi(n)$ such that if ϕ represents a real number x , then the output of $M^\phi(n)$ is

$$M^\phi(n) = \begin{cases} 1 & \text{if } x \in C \\ 0 & \text{if } B(x, 2^{-n}) \cap C = \emptyset \\ 0 \text{ or } 1 & \text{otherwise} \end{cases}$$

It has been shown that the weak definition is equivalent to the standard definition. See [Brv2], for example. We will need the following lemma.

Lemma 3.1. *For any point (z, c) in the complement of the closure $\overline{\mathbb{J}}$, z converges to an attracting periodic orbit of $f_c : z \mapsto z^2 + c$.*

The proof of the lemma occupies §4.

The following lemma allows us to “cover” all points that belong to $\mathbb{J}$.

Lemma 3.2. *There is an algorithm $A_1(n)$ that on input n outputs a sequence of dyadic points $p_1, p_2, \dots \in \mathbb{C} \times \mathbb{C}$ such that*

$$B(\mathbb{J}, 2^{-(n+3)}) \subset \bigcup_{j=1}^{\infty} B(p_j, 2^{-(n+2)}) \subset B(\mathbb{J}, 2^{-(n+1)}).$$

Proof. It is well known that repelling periodic orbits of f_c are dense in J_c . Hence, the set

$$S_{rep} = \{(z, c) : z \text{ is in a repelling periodic orbit of } f_c\}$$

is dense in $\mathbb{J}$. S_{rep} is a union of a countable number of algebraic curves S_{rep}^m given by the constraints

$$\begin{cases} f_c^m(z) = z \\ |(f_c^m)'(z)| > 1 \end{cases}$$

For each m we can compute a finite number of points $p_1^m, \dots, p_{r_m}^m$ approximating S_{rep}^m such that

$$B(S_{rep}^m, 2^{-(n+3)}) \subset \bigcup_{j=1}^{r_m} B(p_j^m, 2^{-(n+2)}) \subset B(S_{rep}^m, 2^{-(n+1)}).$$

We have

$$\overline{\mathbb{J}} = \overline{S_{rep}} = \overline{\bigcup_{m=1}^{\infty} S_{rep}^m}.$$

Hence the computable sequence $p_1^1, \dots, p_{r_1}^1, p_1^2, \dots, p_{r_2}^2, \dots, p_1^m, \dots, p_{r_m}^m, \dots$ satisfies the conditions of the lemma. $\square$

Corollary 3.3. *There is an oracle machine $M_1^{\phi_1, \phi_2}(n)$, where ϕ_1 is an oracle for $z \in \mathbb{C}$ and ϕ_2 is an oracle for $c \in \mathbb{C}$, such that $M_1^{\phi_1, \phi_2}$ always halts whenever $d((z, c), \mathbb{J}) < 2^{-(n+4)}$ and never halts if $d((z, c), \mathbb{J}) \geq 2^{-n}$.*

Proof. Query the oracles for a point $p \in \mathbb{C} \times \mathbb{C}$ such that $d(p, (z, c)) < 2^{-(n+4)}$. Then run the following loop:

$i \leftarrow 0$

do

$i \leftarrow i + 1$

generate p_i using $A_1(n)$ from Lemma 3.2

while $d(p, p_i) > 2^{-(n+2)}$

If $d((z, c), \mathbb{J}) < 2^{-(n+4)}$, then $d(p, \mathbb{J}) < 2^{-(n+3)}$, hence by Lemma 3.2 there is an i such that $d(p, p_i) \leq 2^{-(n+2)}$, and the loop terminates. If $d((z, c), \mathbb{J}) > 2^{-n}$, then $d(p, \mathbb{J}) > 2^{-n} - 2^{-(n+4)} > 1.5 \cdot 2^{-(n+1)}$. Hence, by Lemma 3.2, $p \notin B(p_i, 2^{-(n+1)})$ for all i , and the loop will never terminate. $\square$

The following lemma allows us to exclude points outside $\overline{\mathbb{J}}$ from $\mathbb{J}$.

Lemma 3.4. *There is an oracle machine $M_2^{\phi_1, \phi_2}$, where ϕ_1 is an oracle for $z \in \mathbb{C}$ and ϕ_2 is an oracle for $c \in \mathbb{C}$, such that $M_2^{\phi_1, \phi_2}$ halts if and only if z converges to an attracting periodic orbit (or to ∞) under $f_c : z \mapsto z^2 + c$.*

Proof. M_2 is systematically looking for an attracting cycle of f_c . It also iterates f_c on z with increasing precision and for increasingly many steps until we are sure that either one of the two things holds:

- (1) the orbit of z converges to ∞ ; or
- (2) we find an attracting orbit of f_c and the orbit of z converges to it.

If the search is done systematically, the machine will eventually halt if one of the possibilities above holds. It obviously won't halt if neither holds. $\square$

Proof. (of Theorem 1.5) The algorithm is: Run the machines $M_1^{\phi_1, \phi_2}(n)$ from Corollary 3.3 and $M_2^{\phi_1, \phi_2}$ from Lemma 3.4 in parallel. Output 1 if M_1 terminates first and 0 if M_2 terminates first.

First we observe that $M_1(n)$ only halts on points that are 2^{-n} -close to $\mathbb{J}$, in which case 1 is a valid answer according to Definition 3.1. Similarly, M_2 only halts on points that are outside $\mathbb{J}$, in which case 0 is a valid answer. Hence if the algorithm terminates, it outputs a valid answer. It remains to see that it does always terminate. Consider two cases.

Case 1: $(z, c) \in \mathbb{J}$. In this case $d((z, c), \mathbb{J}) = 0 < 2^{-(n+4)}$, and the first machine is guaranteed to halt.

Case 2: $(z, c) \notin \mathbb{J}$. By Lemma 3.1, z converges to an attracting periodic orbit of f_c in this case, and hence the second machine is guaranteed to halt. $\square$

4. PROOF OF LEMMA 3.1

Suppose $z \notin J_c$ and the orbit of z does not belong to an attracting basin. By the Fatou-Sullivan classification (see e.g. [Mil]), there exists $k \in \mathbb{N}$ such that $w \equiv f_c^k(z)$ belongs to a Siegel disk or to the immediate basin of a parabolic orbit. Our aim is to show that for an arbitrary small $\delta > 0$, there exists a pair $(\tilde{z}, \tilde{c}) \in \mathbb{C} \times \mathbb{C}$ with $|z - \tilde{z}| < \delta$, $|c - \tilde{c}| < \delta$, and for which $\tilde{z} \in J_{\tilde{c}}$. We will treat the Siegel case first.

4.1. The case when w lies in a Siegel disk. Let us denote Δ the Siegel disk containing w , and let $m \in \mathbb{N}$ be its period, that is, the mapping

$$f_c^m : \Delta \rightarrow \Delta$$

is conjugated by a conformal change of coordinates $\phi : \Delta \rightarrow \mathbb{D}$ to an irrational rotation of $\mathbb{D}$.

The following statement is elementary (cf. Prop. 7.1 in [Dou]):

Proposition 4.1. *Denote $\zeta = \phi^{-1}(0) \in \Delta$ the center of the Siegel disk. For each $s > 0$ there exists $\tilde{c} \in B(c, s)$ such that $f_{\tilde{c}}$ has a parabolic periodic point $\tilde{\zeta}$ of period m in $B(\zeta, s)$. In particular, $J_{\tilde{c}}$ is connected, and $B(\zeta, s) \cap J_{\tilde{c}} \neq \emptyset$.*

Consider now the f_c^m -invariant analytic circle

$$S_r = \phi^{-1}(\{z = re^{2\pi i\theta}, \theta \in [0, 2\pi)\})$$

which contains w . Let $\epsilon > 0$ be such that

$$B(w, \epsilon) \subset f_c^k(B(z, \delta)) \cap \Delta.$$

Set $B \equiv B(w, \epsilon/2)$ and let $n \in \mathbb{N}$ be such that the union

$$\bigcup_{0 \leq i \leq n} f_c^{mi}(B) \supset S_r.$$

By Proposition 4.1 for all $\delta > 0$ small enough, there exist $\tilde{c} \in B(c, \delta)$ for which $J_{\tilde{c}}$ is connected and there is a point of $J_{\tilde{c}}$ inside the domain bounded by S_r . Since repelling periodic orbits of f_c are dense in $\partial\Delta$, again for δ small enough, there are points of $J_{\tilde{c}}$ on the outside of S_r as well, and so there exists a point $\xi \in J_{\tilde{c}} \cap S_r$. By construction, there exists $j \in \mathbb{N}$ such that $f_c^j(B(z, \delta)) \ni \xi$. By invariance of Julia set, if $\tilde{c}$ is close enough to c we have $B(z, \delta) \cap J_{\tilde{c}} \neq \emptyset$, and the proof is complete.

4.2. The case when w lies in a parabolic basin. We need to recall the Douady-Lavaurs theory of parabolic implosion ([Dou, Lav]). Denote ζ the parabolic periodic point of f_c whose immediate basin contains w , and let $m \in \mathbb{N}$ be its period.

Recall that an attracting petal P_A is a topological disk whose boundary contains ζ , such that $f_c^{mk}(P_A) \subset P_A$ for some $k \in \mathbb{N}$, and such that the quotient Riemann surface

$$C_A = P_A / f_c^{mk} \simeq \mathbb{C}/\mathbb{Z}.$$

The quotient C_A , called an attracting Fatou cylinder, parametrizes the orbits converging under the dynamics of the iterate f_c^m to ζ inside the periodic cycle of petals $P_A, f_c^m(P_A), \dots, f_c^{m(k-1)}(P_A)$. Recall (see [Mil]) that a quadratic polynomial f_c has only one cycle of petals at ζ . A repelling petal P_R is an attracting petal for the local inverse f_c^{-m} fixing ζ ; the union

$$\bigcup_{0 \leq i \leq k-1} f_c^{mi}(P_A \cup P_R)$$

forms a neighborhood of ζ . The repelling Fatou cylinder C_R is defined in a similar fashion.

Let τ be any conformal isomorphism $C_A \rightarrow C_R$. After uniformization,

$$C_A \xrightarrow{\approx} \mathbb{C}/\mathbb{Z}, \quad C_R \xrightarrow{\approx} \mathbb{C}/\mathbb{Z}$$

$\tau(z) \equiv z + q \pmod{\mathbb{Z}}$ for some $q \in \mathbb{C}$. Let $g : P_A \rightarrow P_R$ be any lift of τ ; it necessarily commutes with f_c^{mk} . Consider the semigroup G generated by the dynamics of the pair (f_c, g) . The orbit Gz of a point $z \in \mathbb{C}$ is independent of the choice of the lift g and only depends on τ .

Set

$$J_{(c, \tau)} = \{z \in \mathbb{C} \text{ such that } Gz \cap J_c \neq \emptyset\}.$$

It can be shown that this set is the boundary of

$$K_{(c, \tau)} = \{z \in \mathbb{C} \text{ such that } Gz \text{ is bounded}\}.$$

The Douady-Lavaurs theory postulates:

Theorem 4.2. *For every τ as above and every $s > 0$ there exists $\tilde{c} \in B(c, s)$ such that $B(J_{\tilde{c}}, s) \supset J_{(c, \tau)}$.*

Since $\zeta \in J_c$, and J_c is connected, there exists a point $u \in J_c \cap P_R$. Let $\hat{w} \in C_A$ be the orbit of w , and let $\hat{u} \in C_R$ be the orbit of u . Choose $\tau : C_A \rightarrow C_R$ so that $\tau(\hat{w}) = \hat{u}$. Then $J_{(c, \tau)} \ni z$, and the claim follows by Theorem 4.2.

REFERENCES

- [BBY1] I. Binder, M. Braverman, M. Yampolsky. Filled Julia sets with empty interior are computable. e-print, math.DS/0410580.
- [BBY2] I. Binder, M. Braverman, M. Yampolsky. On computational complexity of Siegel Julia sets. *Commun. Math. Phys.*, to appear.
- [Brv1] M. Braverman, “Computational Complexity of Euclidean Sets: Hyperbolic Julia Sets are Poly-Time Computable”, Thesis, University of Toronto, 2004, and Proc. CCA 2004, in ENTCS, vol **120**, pp. 17-30.
- [Brv2] M. Braverman, On the Complexity of Real Functions. Proc. of 46th Annual IEEE Symposium FOCS’2005, p. 155-164. Expanded version available as e-print, cs.CC/0502066.
- [Brv3] M. Braverman, Parabolic Julia Sets are Polynomial Time Computable. e-print math.DS/0505036
- [BY] M. Braverman, M. Yampolsky. Non-computable Julia sets. *Journ. Amer. Math. Soc.*, to appear.
- [Dou] A. Douady. Does a Julia set depend continuously on the polynomial? In *Complex dynamical systems*, Proc. Symp. Appl. Math., vol. 49
- [Lav] P. Lavaurs. *Systèmes dynamiques holomorphes, Explosion de points périodiques paraboliques*. Thèse, Orsay, 1989.
- [Mil] J. Milnor. *Dynamics in one complex variable*, 3rd ed. Annals of Math. Studies, Princeton University Press, 2006.
- [RW] R. Rettinger, K. Weihrauch, The Computational Complexity of Some Julia Sets, in STOC’03, June 9-11, 2003, San Diego, California, USA.
- [Wei] K. Weihrauch, *Computable Analysis*, Springer, Berlin, 2000.